



PCT/EP200 4 / 0 1 3 6 9 3 .

18 JAN 2005



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Andrew Gersey

Dated 12 January 2005

BEST AVAILABLE COPY



1/77

THE PATENT OFFICE
LJ

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

RECEIVED BY FAX

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference

044

17NOV03 E852753-1 010198
P01/7700 0.00-0326720.0

2. Patent application number

(The Patent Office will fill in this part)

17 NOV 2003

0326720.0

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

University of Ulster
Office of Innovation and
Enterprise
Science Innovation Centre
Coleraine
BT52 1ST,
N. Ireland, UK

4. Title of the invention

Insulin Releasing Peptides

7054 124003

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

8490567001

Office of Innovation and
Enterprise
Science Innovation Centre
University of Ulster
Coleraine, Co Londonderry
BT52 1ST
N. Ireland, UK.

Patents ADP number (if you know it)

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

YES

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filling with this form. Do not count copies of the same document

Continuation sheets of this form

Description

Claim(s)

Abstract

Drawing(s)

29

26

3

—

—

JF

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

D. Brownlee

Date

17 November 200

12. Name and daytime telephone number of person to contact in the United Kingdom

DAVID BROWNLEE 028 7028 0001

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

"Insulin Releasing Peptides from Skin Secretions of Amphibians"

The present invention relates to the discovery of novel insulin-releasing peptides in the skin secretions of amphibians and their potential use as stimulators of pancreatic beta cell function in the treatment of diabetes mellitus.

Amphibian skin contains two kinds of secretory glands. Mucous glands are distributed throughout the body, and their secretions mainly provide the moist coating necessary for cutaneous respiration. The granular glands, also known as serous or poison glands may be distributed or concentrated in certain areas of the body. The secretions from the granular glands of anurans (frogs and toads) have been shown to contain pharmacologically active substances ranging from simple amines such as norepinephrine and histamine to biologically active peptides, piperidine and steroidal alkaloids, bufadienolides and tetrodotoxin [1-2]. These compounds are thought to play various roles, either in the regulation of physiological functions of the skin or in defence against predators or microorganisms [3-4].

During evolution some of these agents, or closely related daughter molecules, may have taken on distinct physiological roles in humans. It follows that pharmacologically active substances isolated from the skin secretions of frogs and toads may be useful for the treatment of human clinical disease such as diabetes mellitus.

The invention will now be demonstrated with reference to the following non-limiting examples and the accompanying figures wherein:

Figure 1 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Agalychnis calcarifer* stimulate insulin secretion from BRIN-BD11 cells.

Figure 2 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Agalychnis litodryas* stimulate insulin secretion from BRIN-BD11 cells.

Figure 3 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Bombina variegata* stimulate insulin secretion from BRIN-BD11 cells.

Figure 4 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Phyllomedusa trinitatis* stimulate insulin secretion from BRIN-BD11 cells.

Figure 5 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Rana palustris* stimulate insulin secretion from BRIN-BD11 cells.

Figure 6 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Rana pipiens* stimulate insulin secretion from BRIN-BD11 cells.

Figure 7 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Rana saharica* stimulate insulin secretion from BRIN-BD11 cells.

Figure 8 illustrates the dependence of the stimulatory effects of 1653.2 Da purified peptide (peak 1.10) from *Agalychnis calcarifer* on intracellular pathways mediated by protein kinase A (forskolin), protein kinase C (PMA) or pertussis toxin-sensitive G-proteins.

chromatographed on a Vydac 218TP510 semi-preparative C-18 column (25 x 1cm, Hesperia, California, USA). The column was equilibrated with 0.12% (v/v) trifluoroacetic acid/water at a flow rate of 2ml/min. Using 0.1% (v/v) TFA in 70% acetonitrile/water, concentration of acetonitrile in the eluting solvent was raised to 80% (v/v) over 80 min using linear gradients. Absorbance was monitored at 214nm with collection of 2ml fractions. Fractions which showed major insulin releasing activity were pooled and rechromatographed using a Vydac 208TP54 analytical C-18 column (25 x 0.46 cm). The column was equilibrated with 0.12% (v/v) trifluoroacetic acid/water at a flow rate of 1ml/min. Using 0.1% (v/v) TFA in 70% acetonitrile/water, the concentration of acetonitrile in the eluting solvent was raised to 15% (v/v) over 5 min and to 80% (v/v) over 70 min using linear gradients. Absorbance was monitored at 214nm.

Culture of insulin-secreting cells: Clonal rat insulin-secreting BRIN-BD11 cells were cultured in RPMI-1640 tissue culture medium containing 10% (v/v) foetal calf serum, 1% (v/v) antibiotics (100 U/ml penicillin, 0.1 mg/ml streptomycin) and 11.1 mM glucose. The production and characterisation of BRIN-BD11 cells are described elsewhere [5]. Cells were maintained in sterile tissue culture flasks (Corning, Glass Works, UK) at 37 °C in an atmosphere of 5% CO₂ and 95% air using LEEC incubator (Laboratory Technical Engineering, Nottingham, UK). In three experimental series using purified peptides from *Agalychnis calcarifer*, *Bombina variegata* and *Rana sabbavica* cells were cultured overnight with 25 µM forskolin, 10 nM PMA or 0.1 µg/ml pertussis toxin prior to acute tests.

Acute insulin release studies: Insulin release from BRIN-BD11 cells was determined using cell monolayers [5]. The cells were harvested with the aid of trypsin/EDTA (Gibco), seeded into 24-multiwell plates (Nunc, Roskilde, Denmark) at a density of 1.5×10^6 cells per well, and allowed to attach overnight. Prior to acute test, cells were preincubated for 40 min at 37 °C in a 1.0 ml Krebs Ringer bicarbonate buffer (115 mM NaCl, 4.7 mM KCl, 1.28 mM CaCl_2 , 1.2 mM KH_2PO_4 , 1.2 mM MgSO_4 , 10 mM NaHCO_3 , 5 g/l bovine serum albumin, pH 7.4) supplemented with 1.1 mM glucose. Test incubations were performed for 20 min at 37 °C using the same buffer supplemented with 5.6 mM glucose in the absence (control) and presence of various venom fractions, peaks (equivalent to approx. 25 µl dried HPLC fraction) or test agents as indicated in the Figures. Cell viability after 20 min test incubations was assessed by modified neutral red assay [6]. After incubation, aliquots of buffer were removed and stored at -20 °C for insulin radiomimmunoassay [7].

Molecular mass determination: The molecular masses of the purified individual non-toxic peaks exhibiting insulin releasing activity were determined using Matrix Assisted Laser Desorption Ionisation-Time of Flight (MALDI-TOF) mass spectrometry. Electrospray Ionisation quadrupole ion-trap Mass Spectrometry (ESI-MS) was used for *Agalychnis saltator*, *Bombina variegata*, *Rana pipiens* and *Rana saharica*. Masses were recorded and compared with theoretical values calculated by the peptide calculator, a computer software package.

Depyroglutamation: Where necessary (*Bombina variegata*), pyroglutamate at the N-terminal was removed by adding 25 µl of pyroglutamate aminopeptidase preparation (50mM Na_2HPO_4 , 10mM β -mercaptoethanol, 1 mM dithiothreitol, and 1mM EDTA

adjusted to pH 7.3 with H_3HPO_4) containing 0.4 mg/ml pyroglutamate aminopeptidase to 100 μ l of the lyophilised peptide. The reaction mix was incubated for 2 hours at 37 °C and then stored at -20 °C for subsequent amino acid determination by Edman degradation.

Structural analysis by automated Edman degradation: The primary structures of the purified peptides were determined by automated Edman degradation, using an Applied Biosystems Procise 491 microsequencer. Standard operating procedures were used (Applied Biosystems Model 491 Protein Sequencers Users Manual). The limit for detection of phenylthiohydantoin amino acids was 0.2 pmol. The primary structures were compared with those deposited in the SWISSPROT™ database.

Statistical analysis: Results are expressed as mean \pm S.E.M. Values were compared using Student's unpaired t-test. Groups of data were considered to be significantly different if $P < 0.05$.

Results

Isolation, mass spectrometry and sequence analysis of insulin-releasing releasing peptides: Skin secretions from the various amphibian species were purified by HPLC, yielding in each case multiple fractions that were subsequently screened for in vitro biological activity using BRIN-BD11 cells. The insulin-releasing profiles of peaks emerging from the primary HPLC separation are illustrated for *Agalychnis saltator*, *Agalychnis litodryas*, *Bombina variegata*, *Phyllomedusa trinitatis*, *Rana palustris*, *Rana pipiens* and *Rana sylvatica* in Figs. 1-7, respectively.

The major peaks of insulin-releasing activity were subjected to further HPLC purification steps, giving rise ultimately to the isolation of pure peptides with proven insulinotropic activity (Tables 1-7). Where sufficient sample was available, molecular masses and either partial or complete sequences were determined for each peptide as summarised for the various amphibian species in Tables 1-7. In instances where a complete sequence was obtained, the theoretical (calculated) molecular masses of the peptides were shown to corresponded closely to the experimental masses. This indicates the absence of any post-translational modification of constituent amino acids, such as phosphorylation, sulphation or glycation.

Six of the insulinotropic peptides have proved to be established structures. Peak 4.1 from *Rana pipiens* was identical to pipinin-1, peak 23 from *Bombina variegata* to bombesin, peak 2.10 from *Phyllomedusa trinitatis* to dermaseptin BIV precursor and peaks 5.4, 6.5 and 8.3 from *Rana saharica* matched Esculentin-1B, Brevinin-2EC and Brevinin-1E respectively. With these few exceptions, all other insulin-releasing peptides were novel structures as established using the SWISSPROT™ database. Even the functional observations with pipinin-1, dermaseptin BIV precursor, esculentin-1B and brevinin-2EC and 1E were novel as these were totally unsuspected insulin releasing peptides.

Modest similarity existed between some of the isolated insulin-releasing peptides and amphibian antimicrobial peptides that are unsuspected insulin secretagogues such as brevinin, dermaseptin, Rugosin A and tryptophyllin. However, unlike the latter agents, no evidence was obtained of cell lysis or a toxic action that might account for

insulin secretion. Thus the peptides reported herein appear to act through physiological mammalian processes controlling exocytosis of insulin.

To support this view, further studies were carried out using purified peptides from *Agalychnis calcarifer*, *Bombina variegata* and *Rana saharica* to examine cellular mechanisms underlying the stimulation of insulin secretion. The stimulatory effects of the 1653.2 Da peptide (peak 1.10) from *Agalychnis calcarifer* (Figure 8) and 1641.7, 1662.6, 1619.8 and 1650.5 Da peptides (peaks 21, 22, 23 and 24 respectively) from *Bombina variegata* (Figure 9) were abolished in cells cultured overnight with forskolin to desensitise the cyclic AMP-protein kinase A pathway. Overnight culture with PMA or pertussis toxin did not affect the insulin-releasing ability of the peptides, suggesting lack of involvement of protein kinase C or G-protein dependent pathways. Overnight culture with forskolin or PMA resulted in the abolition of the acute stimulatory effects of forskolin or PMA, respectively (Figures 8 and 9). Interestingly, the insulin-releasing action of the 1653.2 Da peptide from *Agalychnis calcarifer* and peaks 21, 22, 23 and 24 from *Bombina variegata* were not affected by 50 μ M verapamil and were clearly evident in cells depolarised with 30 mM KCl (Tables 8 and 9).

The stimulatory effects of the 4920.4 and 4801.2 Da peptides (peaks 5.1 and 5.4) from *Rana saharica* were abolished in cells cultured overnight with forskolin, PMA or pertussis toxin (Figure 10) indicating the involvement of both protein kinase A and C and pertussis toxin-sensitive G-protein in their stimulatory actions. As shown in Table 10, the insulin releasing actions of the isolated peptides were not inhibited by

17-NOV-2003 11:51 AM

the calcium channel blocker verapamil. Stimulatory effects on insulin secretion were also clearly evident in cells depolarised by 30 mM KCl.

Discussion

This research describes for the first time the isolation and characterisation of peptides with insulin-releasing activity from the skin secretions of *Agalychnis calcarifer*, *Agalychnis litodryas*, *Bombina variegata*, *Phyllomedusa trinitatis*, *Rana palustris*, *Rana pipiens* and *Rana saharica*. It is notable that this work has not only uncovered a diverse range of novel peptides structures but it has also revealed that the skin secretions from each amphibian species studied represents an unsuspected and rich source of peptides capable of stimulating physiological insulin secretion from mammalian pancreatic beta cells.

The insulin output induced by amphibian peptides is approximately equivalent to that induced by established mammalian gut peptides, GLP-1, GIP or CCK-8 [8-10]. This indicates that the amphibian peptides isolated are at least as capable as physiological mammalian hormones in stimulating insulin secretion. It is also clear that these peptides may also trigger insulin secretion and have other beneficial actions on beta cells which involve novel secretory pathways as suggested by studies using peak 1.10 from *Agalychnis calcarifer* and peaks 21, 22, 23 and 24 from *Bombina variegata*. In these cases the secretagogues appeared to be mediated through both protein kinase A and G-protein independent pathways. In the case of peptides isolated from *Rana saharica* (peaks 5.1 and 5.4), the stimulatory effects were also independent of pathways triggered by protein kinase C.

It is apparent from the insulin stimulatory effects that specific receptors must exist for these amphibian peptides on mammalian insulin-secreting beta cells. This gives rise to two major and highly novel non-exclusive possibilities. The first is that these insulin-releasing amphibian peptides have homologous or closely related mammalian counter-parts.

The second important possibility arising from this research is that the novel amphibian peptides described in Tables 1-7, or daughter molecules thereof may offer a therapeutically useful means of treating insulin secretory dysfunction and other beta cell disturbances typical of diabetes in humans. Diabetes is predicted to reach epidemic proportions throughout the world in the next 20 years and current treatments do not restore normal glucose homeostasis, therein resulting in debilitating diabetic complications and premature death. Amphibian peptides may therefore be a useful addition to the therapeutic arsenal for use either alone or in combination with other agents to improve diabetes control and decrease the risk of associated complications.

CLAIMS

1. The following peptide structures as stimulators of insulin secretion and pancreatic beta cell function:

- (a) **RRKPLFFIPRPK** (peptide 1.10, *Agalychnis calcarifer*)
- (b) **MLADVFEKIMGD...** (N-terminus of peptide 1.7, *Agalychnis litodryas*)
- (c) **AVWKDFLKNIGKAAGKAVLNSVTDMVNE** (peptide 2.9, *Agalychnis litodryas*)
- (d) **Pyr-QRLGHQWAVGHLM-amidated** (peptide 21, *Bombina variegata*)
- (e) **Pyr-DSFGNQWARCHFM-amidated** (peptide 22, *Bombina variegata*)
- (f) **GKPFYPPPIYPEDM** (peptide 24, *Bombina variegata*)
- (g) **IYNAICPCKHCNKCKPGLLAN** (peptide 25, *Bombina variegata*)
- (h) **XXPLAPFFQAVFK...** (N-terminus of peptide 1.8, *Phyllomedusa trinitatis*)
- (i) **ALWKDILKNVGKAAGKAVLNTVTDMVNQ** (peptide 2.10, *Phyllomedusa trinitatis*)
- (j) **ALSILRGLEKLAKMGIALTNCKATKKC** (peptide 3.8, *Rana palustris*)
- (k) **FLPIAGVAAKVFPKIFCAISKKC** (peptide 4.1, *Rana pipiens*)
- (l) **KGAAGLLEVASCKLSKSC** (peptide 4.22, *Rana saharica*)
- (m) **GIFSKFGRKKIKNLLISGLKNVGKEVGMDVVRTGIDIAGCKIKGEC**
(peptide 5.1 *Rana saharica*)
- (n) **GIFSKLAGKKLKNLLISGLKNVGKEVGMDVVRTGIDIAGCKIKGEC**
(peptide 5.4 *Rana saharica*)
- (o) **GILSTIKDFAIKAGKGAAGLLEMASCKLSGQC** (peptide 5.6, *Rana saharica*)
- (p) **GILLDKLKNFAKTAGKGVLSLLNTASCKLSGQC** (peptide 6.5, *Rana*

saharica)

(g) FLPLLAGLAANFLPKIFCKTRKC (peptide 8.3, Rana saharica)

2. Brevinins, dermaseptins and esculentins as peptides that stimulate insulin secretion by activation of physiological stimulus-secretion coupling pathways rather than by antimicrobial action involving cell lysis.
 3. A peptide as claimed in claims 1 and 2 with at least one amino acid modification by insertion of fatty acid at the alpha amino group of native amino acid or an epsilon amino group of a substituted lysine residue.
 4. A peptide as claimed in claim 1, 2 or 3 having at least one amino acid substitution and/or modification including N-glycated,, N-alkylated, N-acetylated, N-acylated, N-isopropyl, or N-pyroglutamyl amino acids.
 5. Use of a peptide as claimed in any one of claims 1 to 4 in the preparation of a medicament to stimulate insulin secretion and/or moderate blood glucose excursions.
 6. The use of a peptide as claimed in any one of claims 1 to 5 in the preparation of a medicament for treatment of type 1 or type 2 diabetes mellitus.
-
7. A pharmaceutical composition including a peptide according to any one of claims 1 to 6.

8. A pharmaceutical composition useful in the treatment of obesity and/or type 2 diabetes which comprises an effective amount of a peptide as claimed in any of claims 1 to 7 in admixture with a pharmaceutically acceptable excipient for delivery through transdermal, nasal inhalation, oral or injected routes.

9. A pharmaceutical composition as claimed in claim 8 which further comprises one or more sulphonylureas, meglitinides, metformin, and/or thiazolidinediones.

References

- [1] Bevins CL and Zasloff M 1990 Peptides from frog skin. *Annual Review of Biochemistry* 59 395-414
- [2] Erspamer V & Melchiorri P 1983 Neuroendocrine perspectives. pp. 37-104. Elsevier Science, Amsterdam
- [3] Bartholmus GT 1994 Amphibian biology. pp. 382-410. Bcatty and Sons, Surrey.
- [4] Barra D and Simmaco M. 1995 Amphibian skin: a promising resource for antimicrobial peptides. *Trends In Biotechnology* 13 205-209
- [5] McClenaghan NH, Barnett CR, Ah-Sing E, Abdel-Wahab YH, O'Harte FP, Yoon TW, Swanston-Flatt SK and Flatt PR 1996 Characterization of a novel glucose-responsive insulin-secreting cell line, BRIN-BD11, produced by electrofusion. *Diabetes* 45 1132-1140.
- [6] Hunt SM, Chrzanowska C, Barnett CR, Brand HN and Fawell JK 1987 A comparison of in vitro cytotoxicity assays and their application to water samples. *Alternatives to Laboratory Animals* 15 20-29.
- [7] Flatt PR. and Bailey CJ 1981 Abnormal plasma glucose and insulin responses in heterozygous lean (ob/+) mice. *Diabetologia* 20 573-577
- [8] Gault VA, O'Harte FPM, Harriott P, Mooney MH, Green, BD and Flatt PR 2003 Effects of the novel (Pro³) GIP antagonist and exendin (9-39) amide on GIP- and GLP-1-induced cyclic AMP generation, insulin secretion and postprandial insulin release in obese diabetic (ob/ob) mice: evidence that GIP is the major physiological incretin. *Diabetologia* 46 222-230.

- [9] O'Harte FPM, Abdel-Wahab YH, Conlon JM and Flatt PR 1998 Glycation of glucagon-like peptide-1(7-36)amide: characterization and impaired action on rat insulin secreting cells. *Diabetologia* 41 1187-1193.
- [10] Abdel-Wahab YH, O'Harte FPM, Mooney MH, Conlon JM and Flatt PR 1999 N-terminal glycation of cholecystokinin-8 abolishes its insulinotropic action on clonal pancreatic B-cells. *Biochimica et Biophysica Acta* 1452 60-67.

Table 1 Insulin secretion, experimental molecular mass, amino acid sequence, database comparison and theoretical molecular mass of individual peptides isolated from *Agalychnis calcarifer*.

Peptide ID	Insulin release (ng/10 ⁶ cells/20mins)	Experimental Mass (Da)	Amino acid sequence	Database match	Theoretical (Calculated) Mass (Da)
None	1.71 ± 0.12				
1.3	2.52 ± 0.23**	ND	No sequence		
1.10	2.61 ± 0.11**	1653.2	RRKPLFPFIPRPK	No Match	1652.1
1.17	2.10 ± 0.15*	ND	No sequence		
1.18	2.52 ± 0.05***	ND	No sequence		

Table 2 Insulin secretion, experimental molecular mass, amino acid sequence, database comparison and theoretical molecular mass of individual peptides isolated from *Agalychnis litodryas*.

Peptide ID	Insulin release (ng/10 ⁶ cells/20mins)	Experimental Mass (Da)	Amino acid sequence	Database match	Theoretical (Calculated) Mass (Da)
None	1.95 ± 0.16				
1.7	4.22 ± 0.48***	2546.2	MLADVFEKIMGD ...(Insufficient sample)		
2.9	7.46 ± 0.08***	3020.0	AVWKDFLKNIGK AAGKAVLNSVTD MVNE	Dermasept in B IV precursor 79% ID	3019.5

Incubations were performed at 5.6mM glucose. Values are mean ± SEM for 3 separate observations. **P<0.01 and ***P<0.001 compared with 5.6mM glucose. ND

= Not detected. Single letter code denote amino acids: A, Ala; R, Arg; N, Asn; D, Asp; C, Cys; E, Glu; Q, Gln; G, Gly; H, His; X, Hyp; I, Ile; L, Leu; K, Lys; M, Met; F, Phe; P, Pro; S, Ser; T, Thr; W, Trp; Y, Tyr; V, Val.

Table 3 Insulin secretion, experimental molecular mass, amino acid sequence, database comparison and theoretical molecular mass of individual peptides isolated from *Bombina variegata*.

Peptide ID	Insulin release (ng/10 ⁶ cells/20mins)	Experimental Mass (Da)	Amino acid sequence	Database match	Theoretical (Calculated) Mass (Da)
None	1.74 ± 0.08				
21	4.66 ± 0.24***	1641.7	Pyr- QRLGHQWAVG FLM-amidated	Bombesin 93% ID (His) ⁶ Bombesin	1642.7
22	4.75 ± 0.13***	1662.6	Pyr- DSFGNQWARG HFM-amidated	Bombesin 72% ID	1662.9
23	5.67 ± 0.30***	1619.8	Pyr- QRLGNQWAVG FLM-amidated	Bombesin 100% ID	1620.7
24	4.30 ± 0.20***	1650.5	GKPFYPPPIYFE DM	Tryptophyllin 57% ID	1650.9
25	2.39 ± 0.30***	2300.0	IYNAICPCKHCN KCKPGLLAN	No Match	2299.8

Table 4 Insulin secretion, experimental molecular mass, amino acid sequence, database comparison and theoretical molecular mass of individual peptides isolated from *Phyllomedusa trinitatis*.

Peptide ID	Insulin release (ng/10 ⁶ cells/20mins)	Experimental Mass (Da)	Amino acid sequence	Database Match	Theoretical (Calculated) Mass (Da)
None	1.47 ± 0.04				
1.8	2.48 ± 0.37**	8326.4	XXPLAPFFQAVFK ...(Insufficient sample)		
1.11	2.10 ± 0.16**	3379.9	ND		
2.10	2.356 ± 0.34**	2996.4	ALWKDILKNVGKA AGKAVLNTVTDMV NQ	Dermaseptin B IV precursor 100% ID	2998.5

Table 5 Insulin secretion, experimental molecular mass, amino acid sequence, database comparison and theoretical molecular mass of individual peptides isolated from *Rana palustris*.

Peptide ID	Insulin release (ng/10 ⁶ cells/20mins)	Experimental Mass (Da)	Amino acid sequence	Database match	Theoretical (Calculated) Mass (Da)
None	1.467 ± 0.04				
2.6	1.93 ± 0.23**	ND	ND		
2.7	4.14 ± 0.40***	8560.4	ND		
3.1	2.12 ± 0.09***	4919.9	ND		
3.8	2.48 ± 0.44**	2873.5	ALSILRGLEKLAK MGIALTNCKATKK C	Brevinin-1 (46% ID)	2873.7
4.3	2.14 ± 0.13***	3848.7	ND		
4.4	1.87 ± 0.06**	ND	ND		

Table 6 Insulin secretion, experimental molecular mass, amino acid sequence, database comparison and theoretical molecular mass of individual peptides isolated from *Rana pipiens*.

Peptide ID	Insulin release (ng/10 ⁶ cells/20mins)	Experimental Mass (Da)	Amino acid sequence	Database match	Theoretical (Calculated) Mass (Da)
None	2.76 ± 0.13				
3.1	3.46 ± 0.17***	5125.2	ND		
4.1	4.15 ± 0.01***	2562.6	FLPIIAGVAAKV FPKIFCAISKCC	Pipinin-1 100% ID)	2563.2

Incubations were performed at 5.6mM glucose. Values are mean ± SEM for 3 separate observations. **P<0.01 and ***P<0.001 compared with 5.6mM glucose. ND

= Not detected. Single letter code denote amino acids: A, Ala; R, Arg; N, Asn; D,

Asp; C, Cys; E, Glu; Q, Gln; G, Gly; H, His; X, Hyp; I, Ile; L, Leu; K, Lys; M, Met;

F, Phe; P, Pro; S, Ser; T, Thr; W, Trp; Y, Tyr; V, Val

Table 7 Insulin secretion, experimental molecular mass, amino acid sequence, database comparison and theoretical molecular mass of individual peptides isolated from *Rana saharica*.

Peptide ID	Insulin release (ng/10 ⁶ cells/20mins)	Experimental Mass (Da)	Amino acid sequence	Database match	Theoretical (Calculated) Mass (Da)
None	1.87 ± 0.06				
4.14	3.15 ± 0.23**	ND	ND		
4.18	3.52 ± 0.21***	ND	ND		
4.22	3.47 ± 0.40***	1892.6	KGAAGLLEVASC KLSKSC	Rugosin A 68.4% ID	1891.2
4.23	4.25 ± 0.17***	2930.8	AVITGACERDVQC GGGTCCAVSLI.... (insufficient sample)	Protein A/BV8 78% ID	2322.6
4.26	3.08 ± 0.19**	1433.7	ND		
4.27	3.09 ± 0.23**	ND	ND		
4.28	3.19 ± 0.08***	ND	ND		
5.1	3.32 ± 0.22***	4920.4	GIFSKFGRKKIKNL LISGLKNVGKEVG MDVVRTGIDIAGC KIKGEC	Esculentin -1 98% ID	4919.2
5.2	2.95 ± 0.08***	3404.6	ND		
5.3	2.54 ± 0.15**	ND	ND		
5.4	3.30 ± 0.22***	4801.2	GIFSKLAGKKLKN LLISGLKNVGKEV GMDVVRTGIDIAG CKIKGEC	Esculentin -1B 100% ID	4800.8
5.6	2.86 ± 0.37**	3309.2	GILSTKDFAIKAG KGAAGLLEMASC KLSGQC	Brevinin- 2EB 67% ID	3309.0
6.5	5.93 ± 0.47***	3519.3	GILLDKLKNFAKT AGKGVLSLLNTA SCKLSGQC	Brevinin- 2EC 100% ID	3519.2
6.7	3.46 ± 0.28**	3119.2	ND		
8.3	3.53 ± 0.06***	2676.9	FLPLLAGLAANFLP KIFCKTRKC	Brevinin- 1E 100% ID	2676.4

Incubations were performed at 5.6mM glucose. Values are mean ± SEM for 3 separate observations. **P<0.01 and ***P<0.001 compared with 5.6mM glucose. ND = Not detected. Single letter code denote amino acids: A, Ala; R, Arg; N, Asn; D, Asp; C, Cys; E, Glu; Q, Gln; G, Gly; H, His; X, Hyp; I, Ile; L, Leu; K, Lys; M, Met; F, Phe; P, Pro; S, Ser; T, Thr; W, Trp; Y, Tyr; V, Val

Table 8 Effects of the 13 amino acid peptide peak (peak 1.10) from *Agalychnis saltator* on insulin secretion from BRIN-BD11 cells in the presence of verapamil or a depolarising K⁺ concentration.

Addition	Insulin secretion (ng/10 ⁶ cells /20 min)	
	Control	Peak 1.10
None	1.68 ± 0.16	2.75 ± 0.12 *
Verapamil (50μM)	1.74 ± 0.17	3.54 ± 0.20 ΔΔ ***
KCl (30mM)	5.32 ± 0.38 ΔΔΔ	16.45 ± 0.10 ΔΔ ***

Table 9 Effects of peaks 21, 22, 23 and 24 from *Bombina variegata* on insulin secretion from BRIN-BD11 cells in the presence of verapamil or a depolarising K⁺ concentration.

Addition	Insulin secretion (ng/10 ⁶ cells /20 min)				
	Control	Peak 21	Peak 22	Peak 23	Peak 24
None	1.69 ± 0.17	3.65 ± 0.05***	3.23 ± 0.14***	3.40 ± 0.14***	3.31 ± 0.20***
Verapamil (50μM)	1.74 ± 0.17	4.25 ± 0.82***	3.05 ± 0.09***	3.21 ± 0.14***	3.08 ± 0.29***
KCl (30mM)	5.33 ± 0.38 ΔΔΔ	13.68 ± 1.42 ΔΔΔ ***	13.69 ± 1.43 ΔΔΔ ***	15.21 ± 1.43 ΔΔΔ ***	15.35 ± 0.41 ΔΔΔ ***

Acute incubations were performed at 5.6 mM glucose. Values are mean ± SEM for 8 separate observations. *P<0.05 and ***P<0.001 compared with control, ΔΔP<0.01 and ΔΔΔP<0.001 compared with no addition.

Table 10 Effects of peaks 5.1 and 5.4 from *Rana saharica* on insulin secretion from BRIN-BD11 cells in the presence of verapamil or a depolarising K⁺ concentration.

Addition	Insulin secretion (ng/10 ⁶ cells/20 min)		
	Control	Peak 5.1	Peak 5.4
None	1.69 ± 0.17	2.96 ± 0.32***	2.89 ± 0.19***
Verapamil (50μM)	1.74 ± 0.17	2.88 ± 0.28***	2.79 ± 0.10***
KCl (30mM)	5.33 ± 0.38ΔΔΔ	10.11 ± 0.81 ΔΔΔ ***	11.84 ± 0.98 ΔΔΔ ***

Acute incubations were performed at 5.6 mM glucose. Values are mean ± SEM for 8 separate observations. ***P<0.001 compared with control, ΔΔΔP<0.001 compared with no addition.

Legends to Figures

Figure 1 Effects of various semi-preparative C18 HPLC fractions of *Agalychnis calcarifer* crude venom on insulin secretion from BRIN-BD11 cells. Incubations were performed at 5.6mM glucose. Values are the mean ± SEM for 3 separate observations. *P<0.01 and **P<0.001 compared with 5.6mM glucose alone.

Figure 2 Effects of various semi-preparative C18 HPLC fractions of *Agalychnis litodryas* crude venom on insulin secretion from BRIN-BD11 cells. Incubations were performed at 5.6mM glucose. Values are the mean ± SEM for 3 separate observations. *P<0.01 and **P<0.001 compared with 5.6mM glucose alone.

Figure 3 Effects of various semi-preparative C18 HPLC fractions of *Bombina variegata* crude venom on insulin secretion from BRIN-BD11 cells. Incubations were

performed at 5.6mM glucose. Values are the mean \pm SEM for 3 separate observations.
*P<0.01 and **P<0.001 compared with 5.6mM glucose alone.

Figure 4 Effects of various semi-preparative C18 HPLC fractions of *Phyllomedusa trinitatis* crude venom on insulin secretion from BRIN-BD11 cells. Incubations were performed at 5.6mM glucose. Values are the mean \pm SEM for 3 separate observations.
*P<0.001 and **P<0.001 compared with 5.6mM glucose alone.

Figure 5 Effects of various semi-preparative C18 HPLC fractions of *Rana palustris* crude venom on insulin secretion from BRIN-BD11 cells. Incubations were performed at 5.6mM glucose. Values are the mean \pm SEM for 3 separate observations.
*P<0.01 and **P<0.001 compared with 5.6mM glucose alone.

Figure 6 Effects of various semi-preparative C18 HPLC fractions of *Rana pipiens* crude venom on insulin secretion from BRIN-BD11 cells. Incubations were performed at 5.6mM glucose. Values are the mean \pm SEM for 3 separate observations.
*P<0.01 and **P<0.001 compared with 5.6mM glucose alone.

Figure 7 Effects of various semi-preparative C18 HPLC fractions of *Rana saharica* crude venom on insulin secretion from BRIN-BD11 cells. Incubations were performed at 5.6mM glucose. Values are the mean \pm SEM for 3 separate observations.
**P<0.001 compared with 5.6mM glucose alone.

Figure 8 Acute effects of the 13 amino acid 1653.2 Da purified peptide (peak 1.10) from *Agalychnis saltator* on forskolin and PMA on insulin secretion from BRIN-BD11 cells cultured overnight in the absence (control) and presence of 25 μ M forskolin, 10 nM PMA or 0.1 μ g/ml pertussis toxin. Acute incubations were performed

at 5.6 mM glucose. Values are the mean \pm SEM for 8 separate observations.

*** $P < 0.001$ compared with 5.6 mM glucose alone under same culture conditions.

$\Delta\Delta\Delta P < 0.001$ compared with respective test reagent following control culture.

Figure 9 Acute effects of 1641.7, 1662.6, 1619.8 and 1650.5 Da purified peptides (peaks 21, 22, 23 and 24) from *Bombina variegata* on forskolin and PMA on insulin secretion from BRIN-BD11 cells cultured overnight in the absence (control) and presence of 25 μ M forskolin, 10 nM PMA or 0.1 μ g/ml pertussis toxin. Acute incubations were performed at 5.6 mM glucose. Values are the mean \pm SEM for 8 separate observations. ** $P < 0.001$ compared with 5.6 mM glucose alone under same culture conditions. $\Delta P < 0.001$ compared with respective test reagent following control culture.

Figure 10 Acute effects of 4920.4 and 4801.2 Da purified peptides (peaks 5.1 and 5.4) from *Rana saharica* on forskolin and PMA on insulin secretion from BRIN-BD11 cells cultured overnight in the absence (control) and presence of 25 μ M forskolin, 10 nM PMA or 0.1 μ g/ml pertussis toxin. Acute incubations were performed at 5.6 mM glucose. Values are the mean \pm SEM for 8 separate observations. *** $P < 0.001$ compared with 5.6 mM glucose alone under same culture conditions. $\Delta\Delta P < 0.001$ compared with respective test reagent following control culture.

Figure 1 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Agalychnis calcarifer* stimulate insulin secretion from BRIN-BD11 cells.




Figure 2 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Agalychnis litodryas* stimulate insulin secretion from BRIN-BD11 cells.




Figure 3 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Bombina variegata* stimulate insulin secretion from BRIN-BD11 cells.




Figure 4 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Phyllomedusa trinitatis* stimulate insulin secretion from BRIN-BD11 cells.




Figure 5 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Rana palustris* stimulate insulin secretion from BRIN-BD11 cells.




Figure 6 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Rana pipiens* stimulate insulin secretion from BRIN-BD11 cells.




Figure 7 illustrates that semi-preparative C18 HPLC fractions of the skin secretions of *Rana saharica* stimulate insulin secretion from BRIN-BD11 cells.




Figure 8 illustrates the dependence of the stimulatory effects of 1653.2 Da purified peptide (peak 1.10) from *Agalychnis saltator* on intracellular pathways mediated by protein kinase A (forskolin), protein kinase C (PMA) or pertussis toxin-sensitive G-proteins.




Figure 9 illustrates the dependence of the stimulatory effects of 1641.7 Da, 1662.6 Da, 1619.8 Da and 1650.5 Da purified peptides (peaks 21, 22, 23 and 24) from *Bombina variegata* on intracellular pathways mediated by protein kinase A (forskolin), protein kinase C (PMA) or pertussis toxin-sensitive G-proteins.





Figure 10 illustrates the dependence of the stimulatory effects of 4920.4 Da and 4801.2 Da purified peptides (peaks 5.1 and 5.4) from *Rana saharica* on intracellular pathways mediated by protein kinase A (forskolin), protein kinase C (PMA) or pertussis toxin-sensitive G-proteins.



Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP04/013693

International filing date: 17 November 2004 (17.11.2004)

Document type: Certified copy of priority document

Document details: Country/Office: GB
Number: 0326720.0
Filing date: 17 November 2003 (17.11.2003)

Date of receipt at the International Bureau: 10 February 2005 (10.02.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.